

Discovery and quantitative spectral analysis of an Ofpe/WN9 (WN11) star in the Sculptor spiral galaxy NGC 300¹

Fabio Bresolin

Institute for Astronomy, 2680 Woodlawn Drive, Honolulu HI 96822

`bresolin@ifa.hawaii.edu`

Rolf-Peter Kudritzki

Institute for Astronomy, 2680 Woodlawn Drive, Honolulu HI 96822

`kud@ifa.hawaii.edu`

Francisco Najarro

Instituto de Estructura de la Materia, CSIC, Serrano 121, 28006 Madrid, Spain

`najarro@isis.iem.csic.es`

Wolfgang Gieren

Universidad de Concepción, Departamento de Física, Casilla 160-C, Concepción, Chile

`wgieren@coma.cfm.udec.cl`

and

Grzegorz Pietrzyński

*Universidad de Concepción, Departamento de Física, Casilla 160-C, Concepción, Chile and
Warsaw University Observatory, Al. Ujazdowskie 4, 00-478, Warsaw, Poland*

`pietrzyn@hubble.cfm.udec.cl`

ABSTRACT

We have discovered an Ofpe/WN9 (WN11 following Smith et al.) star in the Sculptor spiral galaxy NGC 300, the first object of this class found outside the Local Group, during a recent spectroscopic survey of blue supergiant stars

¹Based on observations obtained at the ESO Very Large Telescope [65.N-0389(B) and 67.D-0006(A)].

obtained at the ESO VLT. The light curve over a five-month period in late 1999 displays a variability at the 0.1 mag level. The intermediate resolution spectra (3800-7200 Å) show a very close resemblance to the Galactic LBV AG Car during minimum. We have performed a detailed non-LTE analysis of the stellar spectrum, and have derived a chemical abundance pattern which includes H, He, C, N, O, Al, Si and Fe, in addition to the stellar and wind parameters. The derived stellar properties and the He and N surface enrichments are consistent with those of other Local Group WN11 stars in the literature, suggesting a similar quiescent or post-LBV evolutionary status.

Subject headings: galaxies: individual (NGC 300) — galaxies: stellar content — stars: Wolf-Rayet

1. Introduction

With the new telescopes of the 8-10 meter class stellar astronomy is branching out beyond the Local Group. Ideal targets for our understanding of young stellar populations in distant galaxies are hot massive stars. These objects have strong stellar winds producing broad and easily detectable spectral features distributed over the whole wavelength range from the UV to the IR, and providing unique information on chemical composition, galactic evolution and extragalactic distances. With these perspectives in mind we have recently begun a systematic spectroscopic study of luminous blue stars in galaxies beyond the Local Group, and presented spectral classification and first quantitative results for A supergiants in NGC 3621 (6.7 Mpc, Bresolin et al. 2001) and NGC 300 (2.0 Mpc, Bresolin et al. 2002). Here we report on the discovery and detailed quantitative analysis of the first example of an Ofpe/WN9 star outside of the Local Group. We will present a detailed chemical abundance pattern – the first in a galaxy beyond the Local Group – together with stellar parameters and a determination of the stellar wind properties.

The Ofpe/WN9 class was introduced to include objects which show in their spectra high excitation emission lines from He II and N III, typical of Of stars, together with low excitation lines from He I and N II, seen in late WN stars (Walborn 1982, Bohannan & Walborn 1989). Objects of this class have so far been identified in the Galaxy (possibly several stars in the Galactic center: Allen et al. 1990, Najarro et al. 1997, Figer et al. 1999), the LMC (ten stars: Bohannan & Walborn 1989), M33 (seven stars: Massey et al. 1996, Crowther et al. 1997) and M31 (one star: Massey & Johnson 1998). The importance of Ofpe/WN9 stars as objects in a transitional stage of evolution between O and W-R stars has been recognized in the last decade, and a connection to the LBV class has been suggested, Ofpe/WN9 stars

being observed during a quiescent or post-LBV phase (Crowther & Smith 1997, Pasquali et al. 1997). Indeed, in at least a couple of instances Ofpe/WN9 stars have been observed to turn into LBVs (R127: Stahl et al. 1983; HDE 269582: Bohannan 1989). The LBV AG Car is also known to show an Ofpe/WN9-like spectrum during its hot phase at visual minimum (Stahl 1986). The discovery of ejected circumstellar nebulae, in some cases measured in a state of expansion, associated with some of the LMC Ofpe/WN9 stars by Nota et al. (1996) and Pasquali et al. (1999) brings forward strong evidence for the occurrence of violent episodes of mass loss in these stars, similar to the shell-producing, eruptive outbursts of LBVs.

Smith et al. (1994) revised and extended the classification of late WN stars to include lower excitation objects. In their scheme, stars showing spectra like AG Car at minimum, where no N III is detected, are reclassified as WN11. Given the spectral resemblance of the NGC 300 star we analyze here to AG Car at visual minimum (see § 4) we will adopt the WN11 classification for the remainder of this Letter. We describe the observational data in § 2, and the photometry in § 3. In § 4 we present our VLT spectra, together with the stellar and wind properties derived from a quantitative spectral analysis.

2. Observations

The data presented here are part of a spectroscopic survey of photometrically selected blue supergiants in the Sculptor spiral galaxy NGC 300 obtained at the VLT with the FORS multiobject spectrograph, and described in detail by Bresolin et al. (2002). This latter paper presents spectra obtained in September 2000 in the blue spectral region ($\sim 4000\text{--}5000\text{ \AA}$) and used for the spectral classification of about 70 supergiants. The emission-line star analyzed here corresponds to star B-16 of the spectral catalog (see Table 2 and finding charts in the aforementioned paper), a rather isolated bright star apparently not associated with any prominent OB association, star cluster or nebulosity. The nearest OB association (from the catalog of Pietrzyński et al. 2001) lies at a deprojected distance of ~ 450 pc. Given the estimated age of 3.8 Myr (§4.1) a speed on the order of 100 km/s would be required for the star to reach its current position from this association. This is much larger than the typical stellar random velocities, and comparable to that of Galactic runaway stars. To the best of our knowledge the star is not included in published listings of emission-line or W-R stars in NGC 300 (Schild & Testor 1992, Breysacher et al. 1997, and references therein), and we will refer to it as star B-16.

In a more recent observing run (September 2001), in order to measure the mass-loss rates of blue supergiants in NGC 300 from the $H\alpha$ line profile, we secured MOS spectra in the red with a 600 lines mm^{-1} grating and $1''$ slitlets, which, together with the older

spectra, provides us with complete coverage at $R \simeq 1,000$ resolution of the 3800-7200 Å wavelength range, except for a 50 Å-wide gap centered at 5025 Å. The total exposure time was 13,500s for both the blue and the red spectra, while better seeing conditions favored the blue observations. The mass-loss rates we derive for the whole blue supergiant sample will be discussed in a forthcoming paper, and we concentrate here on the single B-16 star.

3. Photometry

As part of a multi-epoch photometric study of NGC 300 carried out for the discovery and monitoring of Cepheid variables (Pietrzyński et al. 2001, see also Pietrzyński et al. 2002), BV photometry from ESO/MPI 2.2m telescope WFI images is available for B-16 covering nearly a one-half year period. The resulting B and V light curves are shown in Fig. 1. An apparently irregular variability, on the order of 0.1 mag (peak-to-peak), is detected in both bands, at a significant level above the typical photometric uncertainty of 0.015 mag. The color index remains approximately constant around $B - V = -0.07$. While *bona fide* LBV variables show large amplitude variability on timescales of years (Stahl et al. 2001), partly characterizing the outburst phenomena in these objects, smaller-amplitude variability is observed in Ofpe/WN9 stars or LBVs during quiescence (e.g., Stahl et al. 1984, Sterken et al. 1997). Assuming the intrinsic color index from the stellar atmosphere models ($E_{B-V} = 0.06$), $R = A_V/E_{B-V} = 3.1$ and an average magnitude $V = 19.00$, we obtain from the adopted distance modulus for NGC 300, $m - M = 26.53$ (Freedman et al. 2001), an absolute magnitude $M_V \simeq -7.72 \pm 0.13$.

4. Spectral analysis

The blue and red portions of the rectified VLT optical spectrum are shown in Fig. 2. In the top panel we have also superimposed the spectrum of AG Car during minimum, taken from the Walborn & Fitzpatrick (2000) atlas, in order to illustrate the remarkable similarity between the two spectra. Line identification is provided for most of the recognizable features, which characterize B-16 as a WN11 star. These include the hydrogen Balmer series lines (pure emission), He I lines (together with He II $\lambda 4686$) and mostly low-excitation metal lines such as N II, Fe III and Si II. P-Cygni profiles are seen in most of the He I lines, as well as in Si III $\lambda\lambda 4552$ -4575, N II $\lambda\lambda 4601$ -4643 and Fe III $\lambda\lambda 5127, 5156$. The absence of N III is a discriminant factor against an earlier (WN9-10) classification. The WN11 classification is further confirmed by the observed equivalent widths of He I $\lambda 5876$ (32 Å) and He II $\lambda 4686$ (1 Å), compared to those given by Crowther & Smith (1997).

4.1. Atmosphere models

For the quantitative analysis of the stellar spectrum we have used the iterative, non-LTE line blanketing method presented by Hillier & Miller (1998). The code solves the radiative transfer equation in the co-moving frame for the expanding atmospheres of early-type stars in spherical geometry, subject to the constraints of statistical and radiative equilibrium. Steady state is assumed, and the density structure is set by the mass-loss rate and the velocity field via the equation of continuity. The velocity field (Hillier 1989) is characterized by an isothermal effective scale height in the inner atmosphere, and becomes a β law in the wind (e.g., Lamers et al. 1996). Better fits are obtained if instead of the standard β law a combined 2- β law is used, where each value of β (given in Table 1) characterizes the shape of the velocity field in the inner and outer parts of the wind, respectively. We allow for the presence of clumping via a clumping law characterized by a volume filling factor $f(r)$, so that the ‘smooth’ mass-loss rate, $\dot{M}_S$, is related to the ‘clumped’ mass-loss rate, $\dot{M}_C$, through $\dot{M}_S = \dot{M}_C/f^{1/2}$. The model is then prescribed by the stellar radius R_* , the stellar luminosity L_* , the mass-loss rate $\dot{M}$, the velocity field $v(r)$, the volume filling factor f , and the abundances of the elements considered. The reader is referred to Hillier & Miller (1998, 1999) for a detailed discussion of the code.

The main stellar parameters of our best-fitting model are summarized in Table 1, whereas fits to some of the most important line diagnostics are illustrated in Fig. 3. As can be seen, the majority of the spectral features are well reproduced by the adopted model. The Fe III $\lambda\lambda$ 5127, 5156 lines, together with He II λ 4686, provide excellent constraints for the effective temperature. Using the strength of the electron scattering wings of the Balmer lines we derive a clumping factor $f = 0.25$. Thus the “smooth” mass loss rate would be twice the value displayed in Table 1. The parameters describing the stellar wind, i.e., the terminal velocity $v_\infty = 325 \text{ km s}^{-1}$, the mass-loss rate $\dot{M}_C = 4.6 \times 10^{-5} M_\odot \text{ yr}^{-1}$, and the wind performance number $\eta = \dot{M}v_\infty/(L_*/c) = 0.84$, lie in the range found for Local Group WN11 stars (Crowther et al. 1997). We estimate the uncertainty in L and $\dot{M}_C/f^{1/2}$ to be approximately 10% and 30%, respectively. As a further comparison, Fig. 4 shows the location of star B-16 in the H-R diagram, together with additional WN9-11 stars in the Local Group (Crowther, Hillier & Smith 1995; Smith, Crowther & Willis 1995; Crowther et al. 1997), as well as LBVs at minimum in the Galaxy (P Cyg: Pauldrach & Puls 1990; AG Car: Smith et al. 1994) and in the LMC (R71: Lennon et al. 1994).

Table 2 summarizes the model fractional mass abundances. Uncertainties in the latter range between 0.2 dex (He, N, Si and Fe) and 0.3 dex (C, O and Al). The chemistry of B-16 resembles that of other known Local Group WN9-11 stars. The severe depletion of hydrogen inferred from our analysis (mass fraction $X = 0.27$), and the correspondingly large

helium surface abundance ($H/He = 1.5$ by number), are fairly typical for late WN stars ($H/He = 0.8-3$ by number), and are indicative of heavy mass-loss stripping of the stellar outer layers during the post-main sequence evolution of massive stars. We find a total CNO abundance close to that of Galactic main sequence B stars (Gies & Lambert 1992, Kilian 1992), consistent with a half solar chemical composition in these elements. This agrees with the empirical O abundance derived for an H II region located $30''$ away (region 5 observed by Pagel et al. 1979). On the other hand, Fe has a roughly solar abundance, suggesting an α/Fe ratio different than in the Galaxy. In comparison with the Galactic B star abundance pattern, N is overabundant (by mass) by a factor of 10, while C and O are reduced by factors of ~ 2 and 6, respectively, although the latter abundances are highly uncertain.

The current evolutionary status of B-16 can be estimated by means of the recent Geneva stellar tracks including rotation (Meynet & Maeder 2000) with an initial rotational velocity $v_{ini} = 300 \text{ km s}^{-1}$ and $Z = 0.02$ (Fig. 4). The star’s position in the H-R diagram approaches the $60 M_{\odot}$ track, and an initial mass of $\sim 55 M_{\odot}$ can also be derived from the relation between stellar luminosity and initial mass for WNL stars given by Schaerer & Maeder (1992). From the $60 M_{\odot}$ stellar model more closely matching the observed H and He surface abundances we infer an age of 3.8 Myr and a present mass of $\sim 36 M_{\odot}$. At this stage, the predicted N overabundance with respect to the ZAMS abundance is 12 (mostly occurring during the MS phase, as a consequence of rotational mixing), in good agreement with our finding.

According to its He- and N-enriched surface chemistry, B-16 might be in a dormant or post-LBV phase of evolution. This is also indicated by the position in the H-R diagram, well above the Humphreys-Davidson limit (Humphreys & Davidson 1979), and intermediate between the LBVs AG Car and P Cygni, which have comparable mass-loss rates to B-16, as well as similar wind performance numbers. The NGC 300 star shows higher He enrichment at the surface ($n_{He}/n_H \simeq 0.7$, versus 0.4-0.5 for the LBVs), and a somewhat faster wind (325 versus $185-250 \text{ km s}^{-1}$; Najarro, Hillier & Stahl 1997, Langer et al. 1994), so that a more advanced (post-LBV) state seems more likely. This picture is in agreement with the conclusions reached by Crowther, Hillier & Smith (1995) and Crowther et al. (1997) regarding the close connection between WN9-11 stars and LBVs.

An unexpected reward out of our spectroscopic survey of blue supergiants in NGC 300, the WN11 star B-16 studied here is the first star in this galaxy (and beyond the Local Group) for which we have attempted a detailed quantitative analysis. In the near future the systematic study of the massive stellar population in NGC 300 and similar galaxies within a few Mpc, well within current observational capabilities, will certainly provide new insights into massive stellar evolution and stellar abundances.

We thank J. Hillier for providing his code, and G. Meynet for the Geneva stellar evolutionary tracks including rotation. WG gratefully acknowledges support for this research from the Chilean Center for Astrophysics FONDAF No. 15010003. FN acknowledges Spanish MCYT PANAYA2000-1784 and Ramon y Cajal grants.

REFERENCES

- Allen, D.A., Hyland, A.R., & Hillier, D.J. 1990, MNRAS, 244, 706
- Bohannon, B., & Walborn, N.R. 1989, PASP, 101, 520
- Bohannon, B. 1989, in Physics of Luminous Blue Variables, ed. K. Davidson, A.F.J. Moffat & H.J.G.L.M. Lamers (Dordrecht: Kluwer), p. 35
- Bresolin, F., Gieren, W., Kudritzki, R.-P., Pietrzyński, G., & Przybilla, N. 2002, ApJ, 567, 277
- Bresolin, F., Kudritzki, R.-P., Mendez, R.H., & Przybilla, N. 2001, ApJ, 548, L159
- Breysacher, J., Azzopardi, M., Testor, G., & Muratorio, G. 1997, A&A, 326, 976
- Crowther, P.A., & Smith, L.J. 1997, A&A, 320, 500
- Crowther, P.A., Szeifert, T., Stahl, O., & Zickgraf, F.-J. 1997, A&A, 318, 543
- Crowther, P.A., Hillier, D.J., & Smith, L.J. 1995, A&A, 293, 172
- Figer, D.F., McLean, I.S., & Morris, M. 1999, ApJ, 514, 202
- Freedman, W.L., et al. 2001, ApJ, 553, 47
- Gies, D.R., & Lambert, D.L. 1992, ApJ, 387, 673
- Hillier, D.J., & Miller, D.L. 1999, ApJ, 519, 354
- Hillier, D.J., & Miller, D.L. 1998, ApJ, 496, 407
- Hillier, D.J. 1989, ApJ, 347, 392
- Humphreys, R.M., & Davidson, K. 1979, ApJ, 232, 409
- Kilian, J. 1992, A&A, 262, 171

- Lamers, H.J.G.L.M., Najarro, F., Kudritzki, R.P., Morris, P.W., Voors, R.H.M., van Gent, J.I., Waters, L.B.F.M., de Graauw, T., Beintema, D., Valentijn, E.A., & Hillier, D.J. 1996, *A&A*, 315, 229
- Langer, N., Hamann, W.-R., Lennon, M., Najarro, F., Pauldrach, A.W.A., & Puls, J. 1994, *A&A*, 290, 819
- Lennon, D.J., Wobig, D., Kudritzki, R.-P., & Stahl, O. 1994, in *Evolution of Massive Stars: A Confrontation between Theory and Observation*, eds. D. Vanbeveren, W. van Rensbergen & C. de Loore, *Space Sci. Rev.* 66, 207
- Massey, P., Waterhouse, E., & DeGioia-Eastwood, K. 2001, *AJ*, 119, 2214
- Massey, P., & Johnson, O. 1998, *ApJ*, 505, 793
- Massey, P., Bianchi, L., Hutchings, J.B., & Stecher, T.P. 1996, *ApJ*, 469, 629
- Meynet, G., & Maeder, A. 2000, *A&A*, 361, 101
- Meynet, G., Maeder, A., Schaller, G., Schaerer, D., & Charbonnel, C. 1994, *A&AS*, 103, 97
- Najarro, F., Hillier, D.J., & Stahl, O. 1997, *A&A*, 326, 1117
- Najarro, F., Krabbe, A., Genzel, R., Lutz, D., Kudritzki, R.P., & Hillier, D.J. 1997, *A&A*, 325, 700
- Nota, A., Pasquali, A., Drissen, L., Leitherer, C., Robert, C., Moffat, A.F.J., & Schmutz, W. 1996, *ApJS*, 102, 383
- Pagel, B.E.J., Edmunds, M.G., Blackwell, D.E., Chun, M.S., & Smith, G. 1979, *MNRAS*, 189, 95
- Pasquali, A., Nota, A., & Clampin, M. 1999, *A&A*, 343, 536
- Pasquali, A., Langer, N., Schmutz, W., Leitherer, C., Nota, A., Hubeny, I., & Moffat, F.J. 1997, *ApJ*, 478, 340
- Pauldrach, A.W.A., & Puls, J. 1990, *A&A*, 237, 409
- Pietrzyński, G., Gieren, W., Fouqué, P., & Pont, F. 2002, *AJ*, 123, 789
- Pietrzyński, G., Gieren, W., Fouqué, P., & Pont, F. 2001, *A&A*, 371, 497
- Schaerer, D., & Maeder, A. 1992, *A&A*, 263, 129

- Schild, H., & Testor, G. 1992, A&A, 266, 145
- Smith, L.J., Crowther, P.A., & Willis, A.J. 1995, A&A, 302, 830
- Smith, L.J., Crowther, P.A., & Prinja, R.K. 1994, A&A, 281, 833
- Stahl, O., Jankovics, I., Kovács, J., Wolf, B., Schmutz, W., Kaufer, A., Rivinius, T., & Szeifert, T. 2001, A&A, 375, 54
- Stahl, O. 1986, A&A, 164, 321
- Stahl, O., Wolf, B., Leitherer, C., Zickgraf, F.-J., Krautter, J., & de Groot, M. 1984, A&A, 140, 459
- Stahl, O., Wolf, B., Klare, G., Cassatella, A., Krautter, J., Persi, P., & Ferrari-Toniolo, M. 1983, A&A, 127, 49
- Sterken, C., van Genderen, A.M., & de Groot, M. 1997, in ASP Con. Series 120, Luminous Blue Variables: Massive Stars in Transition, ed. A. Nota & H.J.G.L.M. Lamers (San Francisco: ASP), 35
- Walborn, N.R., & Fitzpatrick, E.L. 2000, PASP, 112, 50
- Walborn, N.R. 1982, ApJ, 256, 45




Fig. 1.— B and V light curves for NGC 300 star B-16, spanning an approximately five month period between September 1999 and January 2000. The $B - V$ color index is shown in the top panel, the V (full dots) and B (open triangles) magnitudes in the bottom panel. Photometric errors (~ 0.015 mag) are comparable to the size of the dots.




Fig. 2.— Rectified VLT/FORS optical spectra obtained at blue (top) and red (bottom) wavelengths. Line identification is provided in the top section of each panel. The comparison spectrum of AG Car at visual minimum (courtesy N. Walborn and E. Fitzpatrick) is shown as a dashed line superimposed on top of the blue spectrum of star B-16.




Fig. 3.— Different spectral diagnostics in the observed spectrum of B-16 compared to our best fit model (dashed line), convolved with a Gaussian profile to match the spectral resolution of the data.




Fig. 4.— Location of B-16 in the H-R diagram (full circle), plotted together with additional WN11 stars (open circles, labeled) and WN9-10 stars (open triangles; from Crowther et al. 1995; Smith et al. 1995; Crowther et al. 1997), as well as LBVs at minimum (open stars, labeled) in the Milky Way (P Cyg: Pauldrach & Puls 1990; AG Car: Smith et al. 1994) and in the LMC (R71: Lennon et al. 1994). We include the Humphreys-Davidson limit (dashed line), and the theoretical Geneva stellar tracks including rotation ($v_{\text{ini}} = 300 \text{ km s}^{-1}$) at solar metallicity, for $M = 40 M_\odot$ and $60 M_\odot$ (dotted lines). These tracks are plotted from the zero-age main sequence to the point where the fractional mass of hydrogen $X \simeq 0.20$.

Table 1. Basic properties and model parameters

R.A. (J2000)	00 54 44.64
Decl. (J2000)	−37 42 39.13
V (mean)	19.00
$B - V$ (mean)	−0.07
M_V	−7.72
<i>Model fit parameters</i>	
E_{B-V}	0.06
$\log(L_*/L_\odot)^a$	5.94
T_* (K) ^a	26000
T_{eff} (K) ^b	22000
$R_*/R_\odot^a$	46
$R_{2/3}/R_\odot^b$	64
$\log(\dot{M}_C) (M_\odot \text{ yr}^{-1})$	−4.34
$v_\infty (km \text{ s}^{-1})$	325
β_1	6.0
β_2	0.9
f	0.25
$\dot{M}_C v_\infty / (L_*/c)$	0.84

^a $\tau \simeq 20$

^b $\tau \simeq 2/3$

Table 2. Model abundances

Species	Relative number fraction	Mass fraction	$X/X_\odot$
H	1.5 E+00	2.7 E−01	0.4
He	1.0 E+00	7.2 E−01	2.7
C	3.6 E−04	7.7 E−04	0.3
N	2.5 E−03	6.3 E−03	7.7
O	5.5 E−04	1.6 E−03	0.2
Al	5.0 E−06	2.4 E−05	0.4
Si	8.5 E−05	1.7 E−04	0.2
Fe	1.5 E−04	1.5 E−03	1.3